



Asymmetric size of ions and orientational ordering of water dipoles in electric double layer model - an analytical mean-field approach



Ekaterina Gongadze, Aleš Iglič*

Laboratory of Biophysics, Faculty of Electrical Engineering, University of Ljubljana, Tržaška 25 and Laboratory of Clinical Biophysics, Faculty of Medicine, Zaloška 9, SI-1000 Ljubljana, Slovenia

ARTICLE INFO

Article history:

Received 27 July 2015

Accepted 30 July 2015

Available online 6 August 2015

Keywords:

Electric double layer
Relative permittivity
Different ions size
Finite volumes
Water dipoles

ABSTRACT

Electric double layer is theoretically described within the mean-field approach by taking into account the orientational ordering of water dipoles and asymmetric size of cations and anions. Analytical expressions for the spatial distribution of ions and water dipoles are derived. The effect of asymmetric ionic size on accumulation of counterions and partial depletion of water molecules near the charged surface, on spatial dependence of relative permittivity and on differential capacitance of electric double layer are presented.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Accumulation of counterions and depletion of coions in the vicinity of a charged surface in contact with an electrolyte solution results in a formation of electric double layer (EDL) [1–15]. Helmholtz initiated the investigation of EDL [16,17] followed by Gouy [18] and Chapman [19] who upgraded the model by considering Boltzmann spatial distribution of the counterions and coions in the Poisson's equation [1,20–22]. Later, Debye and Hückel generalized the Gouy-Chapman model [1,20].

Stern [23] was the first to include the finite size of ions in the EDL model by assuming the distance of closest approach of counterions to the charged surface [24,25]. A more sophisticated approach to take into account the finite size of ions in EDL was introduced by Bikerman [2] and followed by further advance studies [6,8,26–36]. EDL models were further improved by considering the orientational ordering of water dipoles resulting in prediction of local decrease of relative permittivity of electrolyte solution in the saturation regime close to the charged surface [37–39,12,14,15,24,40–44].

Recently, a mean-field model of EDL was developed by Gongadze and Iglič [45] (referred as in this paper as GI model) which encapsulates both, the excluded volume effect (finite size of ions) and orientational ordering of water dipoles, considered as point-like dipoles at the centres of the finite sized spheres with

permittivity equal to the square of optical refractive index of water. In the GI model, the different size of positively and negatively charged ions in EDL was not taken into account [45]. Therefore, in this paper the equations of GI model are generalised in order to incorporate the latter feature. The corresponding analytical expressions for ion spatial distribution functions are derived and included respectively into the Poisson's equation. The influence of different size of ions on the ion and water spatial distribution functions, on spatial dependence of relative permittivity and on differential capacitance for negative and positive voltage are presented and discussed.

2. Theory

2.1. Equal size of ions

By assuming that coions, counterions and water molecules occupy a single lattice site, the number density of water molecules (n_{0w}), counterions (n_0) and coions (n_0) in the bulk electrolyte solution is constant. Therefore, the probability that a single lattice site in a bulk solution is occupied by one of the three kinds of particles (positive ions, negative ions and water molecules) in the electrolyte solution is:

$$\mathcal{P}_+(x \rightarrow \infty) = \mathcal{P}_-(x \rightarrow \infty) = \frac{n_0}{n_0 + n_0 + n_{0w}}, \quad (1)$$

$$\mathcal{P}_w(x \rightarrow \infty) = \frac{n_{0w}}{n_0 + n_0 + n_{0w}}, \quad (2)$$

* Corresponding author.: Tel.: +386 1 4768 825; fax: +386 1 4768 850.
E-mail address: ales.iglic@fe.uni-lj.si (A. Iglič).

The corresponding number densities in a bulk solution are:

$$n_+(x \rightarrow \infty) = n_-(x \rightarrow \infty) = n_s \frac{n_0}{n_0 + n_0 + n_{0w}}, \quad (3)$$

$$n_w(x \rightarrow \infty) = n_s \frac{n_{0w}}{n_0 + n_0 + n_{0w}}, \quad (4)$$

where n_s is the number density of lattice sites:

$$n_s = 2n_0 + n_{0w}. \quad (5)$$

In the vicinity of the charged surface, the number density of ions and water molecules is influenced by the charged surface, so the probabilities that a single lattice site is occupied by a particle of one of the three kinds should be corrected by the corresponding Boltzmann factors, leading to ion and water dipole distribution functions in the form [14,45]:

$$n_+(x) = n_s \frac{n_0 e^{-e_0 \phi \beta}}{n_0 e^{e_0 \phi \beta} + n_0 e^{-e_0 \phi \beta} + n_{0w} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega}, \quad (6)$$

$$n_-(x) = n_s \frac{n_0 e^{e_0 \phi \beta}}{n_0 e^{e_0 \phi \beta} + n_0 e^{-e_0 \phi \beta} + n_{0w} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega}, \quad (7)$$

$$n_w(x) = n_s \frac{n_{0w} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega}{n_0 e^{e_0 \phi \beta} + n_0 e^{-e_0 \phi \beta} + n_{0w} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega}, \quad (8)$$

where $\beta = 1/kT$, kT is the thermal energy, k is the Boltzmann constant, T is the absolute temperature, e_0 is the unit charge and ϕ is the electric potential. Further

$$\begin{aligned} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega &= \frac{2\pi \int_0^\pi d(\cos \omega) e^{-\gamma p_0 E \beta \cos(\omega)}}{4\pi} \\ &= \frac{\sinh(\gamma p_0 E \beta)}{\gamma p_0 E \beta}. \end{aligned} \quad (9)$$

is the water dipole Boltzmann factor after rotational averaging over all possible angles ω [14,45,5]. The angle ω is the angle between the gradient of the electric potential and the vector of the water dipole moment [14,45]. Here p_0 is the magnitude of the external water dipole moment, E is the magnitude of electric field strength. In the model, a single water molecule is considered as a finite sized sphere with permittivity n^2 and a point-like rigid (permanent) dipole/quadrupole at the centre of the sphere, where n is the optical refractive index of water [14,45]. Hence, the cavity field and the electronic polarisability of water molecules are taken into account [46], while the short range interactions between water molecules are neglected. Accordingly, the value of the constant γ is given by [45,14]:

$$\gamma = \frac{3}{2} \left(\frac{2+n^2}{3} \right). \quad (10)$$

2.2. Different size of ions

Consider now a case where negative and positively charged ions occupy more than one lattice site, while a single water molecules still occupy just one lattice site. In the following, α_+ and α_- are the number of lattice sites occupied by a single positive and negative ion, respectively. The probabilities that a single lattice site in the bulk solution is occupied by one of the three kind of particles in electrolyte solution (positive ion, negative ion and water molecule) are then:

$$\mathcal{P}_+(x \rightarrow \infty) = \frac{\alpha_+ n_0}{\alpha_+ n_0 + \alpha_- n_0 + n_{0w}}, \quad (11)$$

$$\mathcal{P}_-(x \rightarrow \infty) = \frac{\alpha_- n_0}{\alpha_+ n_0 + \alpha_- n_0 + n_{0w}}, \quad (12)$$

$$\mathcal{P}_w(x \rightarrow \infty) = \frac{n_{0w}}{\alpha_+ n_0 + \alpha_- n_0 + n_{0w}}, \quad (13)$$

while the corresponding number densities in bulk are

$$n_+(x \rightarrow \infty) = \left(\frac{n_s}{\alpha_+} \right) \frac{\alpha_+ n_0}{\alpha_+ n_0 + \alpha_- n_0 + n_{0w}}, \quad (14)$$

$$n_-(x \rightarrow \infty) = \left(\frac{n_s}{\alpha_-} \right) \frac{\alpha_- n_0}{\alpha_+ n_0 + \alpha_- n_0 + n_{0w}}, \quad (15)$$

$$n_w(x \rightarrow \infty) = n_s \frac{n_{0w}}{\alpha_+ n_0 + \alpha_- n_0 + n_{0w}}, \quad (16)$$

where n_s is the number density of lattice sites:

$$n_s = \alpha_+ n_0 + \alpha_- n_0 + n_{0w}. \quad (17)$$

Similarly as in [14,45], in the vicinity of the charged surface, the number densities of ions and water molecules are corrected by the corresponding Boltzmann factors:

$$n_+(x) = n_s \frac{n_0 e^{-e_0 \phi \beta}}{\alpha_- n_0 e^{e_0 \phi \beta} + \alpha_+ n_0 e^{-e_0 \phi \beta} + n_{0w} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega}, \quad (18)$$

$$n_-(x) = n_s \frac{n_0 e^{e_0 \phi \beta}}{\alpha_- n_0 e^{e_0 \phi \beta} + \alpha_+ n_0 e^{-e_0 \phi \beta} + n_{0w} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega}, \quad (19)$$

$$n_w(x) = n_s \frac{n_{0w} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega}{\alpha_- n_0 e^{e_0 \phi \beta} + \alpha_+ n_0 e^{-e_0 \phi \beta} + n_{0w} \langle e^{-\gamma p_0 E \beta \cos(\omega)} \rangle_\omega}. \quad (20)$$

Taking into account Eq. (9), we can rewrite Eqs. (18)–(20) as:

$$n_+(x) = n_0 e^{-e_0 \phi \beta} \frac{n_s}{\mathcal{D}(\phi, E)}, \quad (21)$$

$$n_-(x) = n_0 e^{e_0 \phi \beta} \frac{n_s}{\mathcal{D}(\phi, E)}, \quad (22)$$

$$n_w(x) = \frac{n_{0w} n_s}{\mathcal{D}(\phi, E)} \frac{\sinh(\gamma p_0 E \beta)}{\gamma p_0 E \beta}. \quad (23)$$

where:

$$\mathcal{D}(\phi, E) = \alpha_- n_0 e^{e_0 \phi \beta} + \alpha_+ n_0 e^{-e_0 \phi \beta} + \frac{n_{0w}}{\gamma p_0 E \beta} \sinh(\gamma p_0 E \beta). \quad (24)$$

The magnitude of the polarisation in the vicinity of the negatively charged surface is given by [14,43,45]:*

$$P(x) = n_w(x) \left(\frac{2+n^2}{3} \right) p_0 \mathcal{L}(\gamma p_0 E \beta). \quad (25)$$

Combining Eq. (23) and Eq. (25) gives the polarization in the form:

$$P(x) = p_0 n_{0w} n_s \left(\frac{2+n^2}{3} \right) \frac{\mathcal{F}(\gamma p_0 E \beta)}{\mathcal{D}(\phi, E)}, \quad (26)$$

where the function $\mathcal{F}(u)$ is defined as: $\mathcal{F}(u) = \mathcal{L}(u)(\sinh u/u)$. Combining $\varepsilon_r(x) = n^2 + P/\varepsilon_0 E$ and Eq. (26) yields the relative (effective) permittivity:

$$\varepsilon_r(x) = n^2 + n_{0w} n_s \frac{p_0}{\varepsilon_0} \left(\frac{2+n^2}{3} \right) \frac{\mathcal{F}(\gamma p_0 E \beta)}{\mathcal{D}(\phi, E) E}. \quad (27)$$

In the limit of vanishing electric field strength ($E \rightarrow 0$) and zero potential ($\phi \rightarrow 0$), the above derived expression for relative permittivity (Eq. (27)) gives the Onsager expression for permittivity [47].

Using the above expression for $\varepsilon_r(x)$, we can rewrite the Poisson's equation into the form of:

$$\frac{d}{dx} \varepsilon_0 \varepsilon_r(x) \frac{d\phi}{dx} = 2 e_0 n_s n_0 \frac{\sinh(e_0 \phi \beta)}{\mathcal{D}(\phi, E)}, \quad (28)$$

* $\mathcal{L}(u)$ is Langevin function

where the macroscopic (net) volume charge density of co-ions and counterions $\rho_{free}(x)$ (Eq. (21) and Eq. (22)) [45] was taken into account:

$$\rho_{free}(x) = e_0 n_+(x) - e_0 n_-(x) = -2 e_0 n_s n_0 \frac{\sinh(e_0 \phi \beta)}{\mathcal{D}(\phi, E)}. \quad (29)$$

The boundary condition at $x=0$ is:

$$\frac{d\phi}{dx}(x=0) = -\frac{\sigma}{\varepsilon_0 \varepsilon_r(x=0)}, \quad (30)$$

where $\varepsilon_r(x)$ is defined by Eq. (27).

3. Results and Discussion

The derived differential equation for spatial distribution of electric potential $\phi(x)$ (Eq. (28)) was solved numerically using Comsol Multiphysics 5.0 by using the boundary condition Eq. (30) at $x=0$.

Fig. 1 shows the spatial dependence of the number density of positive ions (n_+) and water dipoles (n_w) at a negatively charged surface. It can be seen that due to the negative surface charge density σ , the negative ions of the electrolyte solution are nearly completely depleted from the region near the charged surface, while the positively charged ions are accumulated in this region




Fig. 1. The relative number density of positive ions (n_+/n_s) (upper panel) and water dipoles (n_w/n_s) (lower panel) as a function of the distance from a planar charged surface x calculated for bulk concentration of ions $n_0/N_A = 0.1$ mol/l and negative surface charge density $\sigma = -0.2$ As/m² within GI model (full lines) and modified GI model presented in this work (dotted dashed lines). The values of the parameters used are: $\alpha_- = 11$, $\alpha_+ = 5$, dipole moment of water $p_0 = 3.1$ D, optical refractive index $n = 1.33$, bulk concentration of water $n_{ow}/N_A = 55$ mol/l, where N_A is the Avogadro number.




Fig. 2. The relative number density of negative ions (n_-/n_s) (upper panel) and water dipoles (n_w/n_s) (lower panel) as a function of the distance from the planar charged surface (x) calculated for the bulk concentration of ions $n_0/N_A = 0.1$ mol/l and positive surface charge density $\sigma = 0.2$ As/m² within GI model (full lines) and modified GI model (dotted dashed lines). The values of the parameters used here are the same as in Fig. 1.

(Fig. 1). Therefore $\alpha_+ n_+(x=0) + n_w(x=0) \cong n_s$ as it can be seen in Fig. 1.

In contrast to the negatively charged surface (Fig. 1), in the vicinity of the positively charged surface, the negative ions are accumulated and positive ions are completely depleted (Fig. 2). Therefore, in this case $\alpha_- n_-(x=0) + n_w(x=0) \cong n_s$.

A comparison of the results shows that the number density of the positive ions near the negatively charged surface ($n_+(x=0)$ in Fig. 1) is larger than the number density of the negative ions near the positively charged surface ($n_-(x=0)$ in Fig. 2) which can be explained by the larger size of the anions (i.e. $\alpha_- > \alpha_+$). Further, since the anions ($\alpha_- = 11$) are larger than cations ($\alpha_+ = 5$), the number density of water $n_w(x=0)$ at the negatively charged surface (where cations are accumulated) (Fig. 1) is still larger than the number density of water $n_w(x=0)$ at the positively charged surface with accumulated anions of electrolyte solution (Fig. 2). In consequence, the relative permittivity near the charged surface within the modified GI model (presented in this work) is smaller for $\sigma > 0$ than for $\sigma < 0$ as presented in Fig. 3.




Fig. 3. Relative permittivity $\epsilon_r(x)$ as a function of the distance from the charged surfaces at $x=0$ calculated for GI model ($\sigma = -0.2 \text{ As/m}^2$) (full line) and within modified GI model for $\sigma = -0.2 \text{ As/m}^2$ (dotted-dashed line) and $\sigma = 0.2 \text{ As/m}^2$ (dashed line). The values of the other model parameters are the same as in Fig. 1.

It has been shown that the calculated differential capacitance of the EDL within GC model monotonously increases as a function of the increasing surface potential, while in Bikerman [10,12,48] and GI models [14] starts to decrease after reaching a maximal value, as shown in Fig. 4. For higher values of the surface potential, the calculated GI differential capacitance drops below the corresponding Bikerman values to very small values [14] in accordance with the experimental results [49,50]. As presented in Fig. 4, the differential capacitance decreases even more within the modified GI model. As expected, consideration of different sizes of positive and negative ions within modified GI model may describe some characteristics of the experimentally observed asymmetric camel-like dependence of differential capacitance on surface potential/voltage [49–51]. The latter was already studied within an approximate approach for small volume shares of ions everywhere in the electrolyte solution, where the number densities of ions and water molecules were obtained implicitly [52]. In contrast, here the number densities of ions and water molecules are given explicitly by analytic expressions (Eqs. (21)–(23)). Furthermore, the assumption of small volume shares of ions everywhere in the electrolyte solution was not applied here because for larger magnitudes of surface charge density, the volume share of counterions near the charged surface is in general not small.




Fig. 4. Differential capacitance as a function of the surface potential ϕ_0 calculated within GI and modified GI model for bulk concentration of ions $n_0/N_A = 0.1 \text{ mol/l}$. The values of other model parameters are given in Fig. 1.

Further generalisation of the modified GI given in this paper model should include also the Helmholtz/Stern layer taking into account unequal distances of closest approach for cations and anions ions [25,53–55]. This modification would change the relative height for both maxima of the differential capacitance camel-like curve [25] and also move them to higher magnitudes of the potential/voltage [25,56].

4. Conclusion

In this work, the asymmetric size of cations and anions in the electric double layer are described within a mean-field approach and lattice statistics as a generalisation of our previous published GI model [45], which takes into account spatial dependence of relative permittivity due to orientational ordering of water dipoles in saturation regime and partial depletion of water near the charged surface [14]. Within the presented modified GI model, we derived analytical expressions for the spatial distribution of ions and water dipoles in the electric double layer. As illustration, it is shown that the sign of the surface charge of the planar charged surface influences partial depletion of water molecules and local decrease of relative permittivity near the charged surface. The influence of size asymmetry of cations and anions in electrolyte solution on differential capacitance of EDL is also investigated.

For real systems our model can be applied to biological systems where the cell and liposome membranes are in contact with electrolyte solution, usually with predominant sodium cations and chloride anions [3–5,21]. The model can be also used to study the cell membrane interactions with different biomaterials/implants or interaction of cells with inorganic nanoparticles and nanostructured surfaces [14]. In colloid science [22] and electrochemistry [1,24], the presented model can be used in the studies of electrode/ionic liquids interface, as for example in calculation of differential capacitance [24,10,12,48,14,52] (Fig. 4) where the metallic or semiconductor electrodes are in contact with different kind of electrolytes having asymmetric size of ions [51]. Recently, hybrid nanoporous semiconductor electrodes [57], covered by organic materials, are gaining more and more interest, as for example in construction of supercapacitors [58].

Acknowledgements

This work was supported by Slovenian Research Agency (ARRS) grant P2-0232.

References

- [1] E. Verwey, J. Overbeek, *Theory of the Stability of Lyophobic Colloids*, Dover Publications, Inc, Mineola, New York, 1999.
- [2] J. Bikerman, Structure and capacity of electrical double layer, *Phil. Mag.* 33 (1942) 384–397.
- [3] S. McLaughlin, The electrostatic properties of membranes, *Annual Review of Biophysics and Biophysical Chemistry* 18 (1) (1989) 113–136.
- [4] G. Cevc, Membrane electrostatics, *Biochim. Biophys. Acta* 1031 (1990) 311–382.
- [5] A. Safran, *Statistical Thermodynamics of Surfaces, Interfaces, and Membranes*, Addison-Wesley Publishing Company, 1994.
- [6] E. Wicke, M. Eigen, Über den Einfluss des Raumbedarfs von Ionen in wässriger Lösung auf ihre Verteilung im elektrischen Feld und ihre Aktivitätskoeffizienten, *Z. Elektrochem.* 56 (1952) 551–561.
- [7] R. Netz, Electrostatics of counter-ions at and between planar charged walls: From Poisson-Boltzmann to the strong-coupling theory, *Eur. Phys. J. E* 5 (2001) 557–574.
- [8] S. Lamperski, C.W. Outhwaite, Exclusion volume term in the inhomogeneous Poisson-Boltzmann theory for high surface charge, *Langmuir* 18 (2002) 3423–3424.
- [9] I. Bivas, Electrostatic and mechanical properties of a flat lipid bilayer containing ionic lipids, *Coll. Surf. A* 282 (283) (2006) 423–434.
- [10] A. Kornyshev, Double-layer in ionic liquids: paradigm change? *Chem. Phys. Lett.* 111 (2007) 5545–5557.

- [11] G. Tresset, Generalized Poisson-Fermi formalism for investigating size correlation effects with multiple ions, *Phys. Rev. E* 78 (2008) 061506.
- [12] M. Bazant, M. Kilić, B. Storey, A. Ajdari, Towards an understanding of induced-charge electrokinetics at large applied voltages in concentrated solutions, *Adv. Coll. Int. Sci.* 152 (2009) 48–88.
- [13] Z. Arsov, M. Rappolt, J. Grdodolnik, Weakened hydrogen bonds in water confined between lipid bilayers: the existence of a long-range attractive hydration force, *Chem. Phys. Chem.* 10 (9) (2009) 1438–1441.
- [14] E. Gongadze, A. Velikonja, Š. Perutkova, P. Kramar, A. Maček-Lebar, V. Kralj-Iglič, A. Igljič, Ions and water molecules in an electrolyte solution in contact with charged and dipolar surfaces, *Electrochim. Acta* 126 (2014) 42–60.
- [15] M.A. Quiroga, K.H. Xue, T.K. Nguyen, M. Tulodziecki, H. Huang, A.A. Franco, A multiscale model of electrochemical double layers in energy conversion and storage devices, *J. Electrochem. Soc.* 161 (2014) E3302–E3310.
- [16] H. Helmholtz, Über einige Gesetze der Verteilung elektrischer Ströme in körperlichen Leitern, mit Anwendung auf die thierisch-elektrischen Versuche, *Ann. Phys. Chem.* 165 (6) (1853) 211–233.
- [17] H. Helmholtz, Studien über elektrische Grenzschichten, *Ann. Phys. Chem.* 243 (1879) 337–382.
- [18] G. Gouy, Sur la constitution de la charge électrique à la surface d'un électrolyte, *J. Phys. Radium* 9 (1910) 457–468.
- [19] D. Chapman, A contribution to the theory of electrocapillarity, *Philos. Mag.* 25 (1913) 475–481.
- [20] F. Fogolari, A. Brigo, H. Molinari, The Poisson-Boltzmann equation for biomolecular electrostatics: a tool for structural biology, *J. Mol. Recognit.* 15 (6) (2002) 377–392.
- [21] D. Andelman, *Handbook of Biological Physics, Vol 1*, Elsevier Science B.V., 1995, Ch. Electrostatics properties of membranes: The Poisson-Boltzmann theory, pp. 603–642.
- [22] D.F. Evans, H. Wennerström, *The Colloidal Domain: where physics, chemistry, biology, and technology meet*, 2nd ed., Wiley-VCH, New York, 1999.
- [23] O. Stern, Zur Theorie der elektrolytischen Doppelschicht, *Z. Elektrochem.* 30 (1924) 508–516.
- [24] H.J. Butt, K. Graf, M. Kappl, *Physics and Chemistry of Interfaces*, Wiley-VCH, Weinheim, 2003.
- [25] A. Velikonja, V. Kralj-Iglič, A. Igljič, On asymmetric shape of electric double layer capacitance curve, *Int. J. Electrochem. Sci.* 10 (2015) 1–7.
- [26] L. Bhuiyan, C. Outhwaite, D. Bratko, Structure and thermodynamics of micellar solutions in the modified Poisson-Boltzmann theory, *Chem. Phys. Lett.* 193 (1992) 203–210.
- [27] S.W. Kenkel, J.R. Macdonald, A lattice model for the electrical double layer using finite-length dipoles, *J. Chem. Phys.* 81 (1984) 3215–3222.
- [28] P. Nielaba, F. Forstmann, Packing of ions near an electrolyte-electrode interface in the hnc/lmsa approximation to the rpm model, *Chem. Phys. Lett.* 117 (1) (1985) 46–48.
- [29] C. Caccamo, G. Pizzimenti, L. Blum, An improved closure for the Born-Green-Yvon equation for the electric double layer, *J. Chem. Phys.* 84 (6) (1986) 3327–3335.
- [30] R. Kjellander, S. Marčelja, Interaction of charged surfaces in electrolyte solutions, *Chem. Phys. Lett.* 127 (4) (1986) 402–407.
- [31] M. Plischke, D. Henderson, Pair correlation functions and density profiles in the primitive model of the electric double layer, *J. Chem. Phys.* 88 (4) (1988) 2712–2718.
- [32] L. Mier-y Teran, S. Suh, H.S. White, H. Davis, A nonlocal free-energy density-functional approximation for the electrical double layer, *J. Chem. Phys.* 92 (8) (1990) 5087–5098.
- [33] P. Strating, F. Wiegand, Effects of excluded volume on the electrolyte distribution around a charged sphere, *J. Phys. A Math. Gen.* 26 (1993) 3383–3391.
- [34] V. Kralj-Iglič, A. Igljič, A simple statistical mechanical approach to the free energy of the electric double layer including the excluded volume effect, *J. Phys. II France* 6 (1996) 477–491.
- [35] I. Borukhov, Charge renormalization of cylinders and spheres: ion size effect, *J. Pol. Sci. B, Pol. Phys.* 42 (2004) 3598–3615.
- [36] P. Biesheuvel, M. van Soestbergen, Counterion volume effects in mixed electrical double layers, *J. Coll. Int. Sci.* 316 (2007) 490–499.
- [37] C.W. Outhwaite, A treatment of solvent effect in the potential theory of electrolyte solution, *Mol. Phys.* 31 (5) (1976) 1345–1357.
- [38] C.W. Outhwaite, Towards a mean electrostatic potential treatment of an ion-dipole mixture or a dipolar system next to a plane wall, *Mol. Phys.* 48 (3) (1983) 599–614.
- [39] A. Abrashkin, D. Andelman, H. Orland, Dipolar Poisson-Boltzmann equation: ions and dipoles close to charge interfaces, *Phys. Rev. Lett.* 99 (2007) 077801.
- [40] E. Gongadze, T. Slivnik, V. Kralj-Iglič, A. Igljič, The quadrupole moment of water molecules and the permittivity of water near a charged surface, *Electrochim. Acta* 109 (2013) 656–662.
- [41] T. Nagy, D. Henderson, D. Boda, Simulation of an electrical double layer model with a low dielectric layer between the electrode and the electrolyte, *J. Phys. Chem. B* (2011) 11409–11419.
- [42] A. Igljič, E. Gongadze, K. Bohinc, Excluded volume effect and orientational ordering near charged surface in solution of ions and Langevin dipoles, *Bioelectrochemistry* 79 (2010) 223–227.
- [43] E. Gongadze, U. van Rienen, V. Kralj-Iglič, A. Igljič, Langevin Poisson-Boltzmann equation: point-like ions and water dipoles near a charged membrane surface, *Gen. Physiol. Biophys.* 30 (2) (2011) 130–137.
- [44] R. Misra, S. Das, S. Mitra, Electric double layer force between charged surfaces: Effect of solvent polarization, *J. Chem. Phys.* 138 (2013) 114703.
- [45] E. Gongadze, A. Igljič, Decrease of permittivity of an electrolyte solution near a charged surface due to saturation and excluded volume effects, *Bioelectrochemistry* 87 (2012) 199–203.
- [46] H. Fröhlich, *Theory of dielectrics*, Clarendon Press, Oxford, UK, 1964.
- [47] L. Onsager, Electric moments of molecules in liquids, *J. Am. Chem. Soc.* 58 (1936) 1486–1493.
- [48] M.V. Fedorov, N. Georgi, A. Kornyshev, Double layer in ionic liquids: The nature of the camel shape of capacitance, *Electrochem. Commun.* 12 (2010) 296–299.
- [49] V. Lockett, R. Sedev, J. Ralston, M. Horne, T. Rodopoulos, Differential capacitance of the electrical double layer in imidazolium based ionic liquids - influence of potential, cation size, and temperature, *J. Phys. Chem. C* 1124 (2008) 7486–7495.
- [50] V. Lockett, M. Horne, R. Sedev, T. Rodopoulos, J. Ralston, Differential capacitance of the double layer at the electrode/ionic liquids interface, *Phys. Chem. Chem. Phys.* 12 (2010) 12499–12512.
- [51] D. Grahame, Differential capacity of mercury in aqueous sodium fluoride solutions. I. effect of concentration at 25 degrees, *J. Am. Chem. Soc.* 76 (1954) 4819–4823.
- [52] J. Sin, S. Im, K. Kim, Asymmetric electrostatic properties of an electric double layer: a generalized Poisson-Boltzmann approach taking into account non-uniform size effects and water polarization, *Electrochim. Acta* 153 (2015) 531–539.
- [53] J. Yu, G. Aguilar-Pineda, A. Antillon, S. Dong, M. Lozada-Cassou, The effects of unequal ionic sizes for an electrolyte in a charged slit, *J. Colloid Interface Sci.* 295 (2006) 124–134.
- [54] C. Outhwaite, L. Bhuiyan, A modified Poisson-Boltzmann equation in electric double layer theory for a primitive model electrolyte with size asymmetric ion size, *J. Chem. Phys.* 84 (1986) 3461–3471.
- [55] E. Gongadze, A. Igljič, Relative permittivity in Stern and diffuse layers, *Acta Chim. Slov.* 61 (2014) 241–245.
- [56] A. Velikonja, E. Gongadze, V. Kralj-Iglič, A. Igljič, Charge dependent capacitance of Stern layer and capacitance of electrode/electrolyte interface, *Int. J. Electrochem. Sci.* 9 (2014) 5885–5894.
- [57] F. Fabregat-Santiago, G. Garcia-Belmonte, J. Bisquert, P. Bogdanoff, A. Zaban, Mott-Schottky analysis of nanoporous semiconductor electrodes in dielectric state deposited on SnO₂(F) conducting substrates, *J. Electrochem. Soc.* 150 (2003) E293–E298.
- [58] R. Imani, M. Pazoki, A. Tiwari, G. Boschloo, A. Turner, V. Kralj-Iglič, A. Igljič, Band edge engineering of TiO₂@DNA nanohybrids and implications for capacitive energy storage devices, *Nanoscale* 7 (2015) 10438–10448.